

RX

Drug Name: Epinephrine
Trade Name: Adrenalin, Epi

REVISED: November 1, 2019

Class:

- Adrenergic Catecholamine
- Sympathomimetic

Mechanism of Action:

- β_1 —increases contractility (positive inotrope), AV conduction (positive dromotrope), and automaticity
- β_2 --bronchodilation, skeletal muscle vasodilation
- α --peripheral vasoconstriction, fight or flight response
- Small doses, beta effects dominate--vasodilation
- Large doses, alpha effects dominate--vasoconstriction, increases systemic vascular resistance and blood pressure

Indications:

- Hypersensitivity reactions (anaphylaxis)
- Acute bronchospasm associated with asthma or COPD (refractory to first-line agents)
- Asystole, VF, pulseless VT, PEA
- Croup & epiglottitis

Contraindications:

- None in cardiac arrest or severe anaphylaxis
- Hypersensitivity

Precautions:

- HTN
- Ischemic heart disease
- Cerebrovascular insufficiency
- Deactivated/precipitates with alkaline solutions (NaHCO₃)
- Increases myocardial oxygen demand
- Pulmonary edema
- Pregnancy (C)
- Geriatrics
- Protect from light

Dosage:

Adults:

Pulseless Rhythms

- **IV/IO:** 1 mg (1:10,000) every 3-5 minutes

Anaphylaxis

- **IM/SQ:** 0.3-0.5 mg (1:1,000), repeat once at 10 minutes if s/s do not improve
- **IV Infusion:** IV/IO: 0.05-1 mcg/kg/min titrate for effect
 - **For refractory Cases**
 - **To Mix:** 1 mg epinephrine in 250 cc NS bag
- **Neb:** *For laryngeal edema only*, 3 mg epinephrine 1:1,000 (3 ml) mixed with 3 ml NS for 6ml solution total

DRUG: EPINEPHRINE

RX

This document is for **reference only**. Please refer to SWO's for specific indications, dosages, and applications

DRUG: EPINEPHRINE

Acute bronchospasm associated with asthma or COPD (refractory to first-line agents)

- **IM/SQ:** 0.3-0.5 mg (1:1,000)

Persistent/Refractory Hypotension

- **IV Infusion:** IV/IO: 0.05-1 mcg/kg/min , titrate for effect
 - **For refractory Cases**
 - **To Mix:** 1 mg epinephrine in 250 cc NS bag

Symptomatic Ca Channel Blocker/Beta Blocker OD refractory to other interventions

- **IV Infusion: IV/IO: 0.05-1 mcg/kg/min titrate for effect**
 - **For refractory Cases**
 - **To Mix:** 1 mg epinephrine in 250 cc NS bag

Pediatrics:

Pulseless Rhythms:

- **IV/IO:** 0.01 mg/kg (1:10,000) every 3-5 minutes
- **NEONATES:** 0.01-0.03 mg/kg (1:10,000) IV/IO every 3-5 minutes

Anaphylaxis

- **IM/SQ:** 0.01 mg/kg (1:1,000), **MAX: 0.3 mg**
- **Neb:** *For laryngeal edema only*, 3 mg epinephrine 1:1,000 (3 ml) mixed with 3 ml NS for 6ml solution total

Persistent/Refractory Hypotension

- **IV Infusion:** 0.05-1 mcg/kg/min, titrate for effect
 - **To Mix:** 1 mg epinephrine in 250 cc NS bag

Croup & Epiglottitis:

- **Neb:** *For laryngeal edema only*, 3 mg epinephrine 1:1,000 (3 ml) mixed with 3 ml NS for 6ml solution total

Refractory Bronchospasm (Severe):

- **IM/SQ:** 0.01 mg/kg (1:1000, 0.1 ml/kg)

Onset:

- IV/IO: 1-2 min
- IM/SQ: 5-10 min

Duration:

- IV/IM/SQ: 5-10 min

Side Effects:

- Anxiety
- Tachycardia
- HTN
- Angina
- Arrhythmias
- V-Fib
- N/V
- Fear
- Headache
- Pallor
- Dizziness
- Tremors

This document is for **reference only**. Please refer to SWO's for specific indications, dosages, and applications

RX

Interactions:

- Potentiated by MAOIs and TCAs
- Antagonized by beta blockers
- Precipitates in alkaline solutions

PEARLS:

CAUTION: All patients receiving inhaled beta agonists and/or anticholinergic medications should be observed for a least one hour following treatment for return of symptoms.

ALS evaluation is indicated if Epi administered either PTA or by EMS, and transport strongly encouraged. Refusals require medical control contact.

- **I.M. Epi may be more effective than SQ Epi in shock situations.**
- Sodium bicarbonate or Furosemide will inactivate epinephrine; flush line well between administration.
- Use an IV Infusion pump when administering Epi Infusions.
 - **To Mix:** 1 mg epinephrine in 250 cc NS bag

DRUG: EPINEPHRINE

RX

BLANK PAGE

DRUG: EPINEPHRINE

REFERENCE ONLY