

RX

Drug Name: **Lidocaine Hydrochloride**
Trade Name: **Lidocaine, Xylocaine**
REVISED: **November 1, 2018**

Class:

- Antidysrhythmic (Class I-B)

Mechanism of Action:

- Decreases ventricular automaticity (*reduces the slope of phase 4 diastolic depolarization*)
- Decreases excitability, and raises fibrillation threshold
- Decreases conduction in ischemic cardiac tissue without adversely affecting normal conduction
- Does not affect contractility

Indications:

- Pulseless Ventricular Tachycardia, Ventricular Fibrillation.
- Ventricular Tachycardia with a pulse
- Malignant PVCs
- Wide complex tachycardias of unknown origin
- Pre-Intubation in the setting of closed head injuries and strokes

Contraindications:

- Advanced AV block (2nd degree Type II (Mobitz II) and 3rd degree blocks) in the absence of a functioning pacemaker
- Adams-Stokes syndrome

Precautions:

- Hypotension
- Torsades de Pointes (*if known torsade's, magnesium is the agent of choice*)

Dosage:

Adults:

- **Pulseless VT, VF:** IV/IO: 1.0 to 1.5 mg/kg IV bolus, can repeat in 3-5 minutes **not to exceed 3 mg/kg or 300 mg in 30 minutes (not including infusion)**
- **Ventricular ectopy, and Wide complex tachycardia:** 1.0-1.5 mg/kg slow IV bolus followed by additional doses of 0.5-0.75 mg/kg every 5minutes **not to exceed 3 mg/kg or 300 mg in 30 minutes (not including infusion)**. If ectopy resolves, can set up a continuous infusion. (Be sure to rebolus @ 0.5-0.75 mg/kg in 8-10 minutes to maintain therapeutic levels of lidocaine)
- **Continuous infusion:** 2-4 mg/minute titrated for effect, to be initiated if V-fib/V-Tach resolves. (Start @ 2 mg/min & add 1 mg/min for each additional 1 mg/kg IV bolus)
 - 1 mg/kg bolus = 2 mg/min.
 - 1.5-2 mg/kg total bolus = 3 mg/min.
 - 2.5-3 mg/kg total bolus = 4 mg/min.
- Management of Pain with Intraosseous infusions
 - IO: 20-40 mg administered slow every 30 minutes **not to exceed 3 mg/kg or 300 mg**

DRUG: Lidocaine

RX

This document is for **reference only**. Please refer to SWO's for specific indications, dosages, and applications

DRUG: Lidocaine

Pediatrics:

- All Ventricular dysrhythmias:
 - IV/IO: 1 mg/kg bolus, not to exceed 3 mg/kg
- Continuous Infusion
 - IV/IO: 20-50 $\mu\text{g/kg/min}$ infusion. 120 mg in 100 ml NS yielding 1.2 mg/ml.
 - *If a bolus dose of lidocaine has not been administered within the previous 15 minutes, administer a bolus of 1 mg/kg before initiation of the infusion*
- Management of Pain with Intraosseous infusions
 - IO: 0.5 mg/kg administered slow every 30 minutes **not to exceed 3 mg/kg**
 - MAX single dose 40 mg.

Onset:

IV—30-90 sec

Duration:

IV—10-20 minutes for bolus

Side Effects:

CNS--drowsiness, dizziness, confusion, slurred speech, seizures, respiratory depression/arrest
CV--hypotension, dysrhythmias, bradycardia, cardiac arrest
Other--Nausea, vomiting

Interactions:

Additive cardiac depression and toxicity when used concomitantly with amiodarone, procainamide, phenytoin, quinidine, & beta-blockers

PEARLS:

- Evidence for one particular antiarrhythmic over another is inconclusive.
- Previously, lidocaine was used for pre-treatment of increased inter-cranial pressure during RSI. A review of the evidence has shown no clear benefit of this practice, and an increased risk of hypotension which can be detrimental in the neuro-critically ill. **Therefore the use of Lidocaine during RSI for increased ICP is no longer recommended.**
- Always give full initial dose, but reduce all subsequent doses by $\frac{1}{2}$ for elderly (>70) or with impaired hepatic function
- Lidocaine has been shown to prolong apnea when used in conjunction with neuromuscular blocking agents. This usually occurs at higher doses than the dose used for pre-intubation, but be aware
- If you have a bradycardia with PVCs, always treat the bradycardia first (atropine, pacing, etc.), prior to threatening the PVCs

REFERENCE ONLY

- Lidocaine jelly works well as a topical anesthetic for intubation, use it to lubricate ETT tubes, but not the stylette as the viscosity of lidocaine jelly may occlude your tube
- Lidocaine induced CNS toxicity usually presents with clinical signs of CNS stimulation followed by CNS depression. Cardiac toxicity follows CNS toxicity and usually requires very high doses

RX

DRUG: Lidocaine

RX

DRUG: Lidocaine

REFERENCE ONLY